

**After Immunization Information and Record – Please Keep!**

Your child \_\_\_\_\_ has received the following immunization(s):

- Diphtheria, tetanus, acellular pertussis and inactivated polio(DTaP-IPV/Tdap-IPV)** Date: \_\_\_\_\_  
**Possible side effects:** Soreness, swelling and/or redness at the injection site; a small painless lump at the injection site that may take a few weeks to disappear but does not need any treatment; fever (temperature of 38.5 C or higher); muscle or joint pain; tiredness or drowsiness; irritability or crying; vomiting; may feel generally unwell.
  
- Meningococcal conjugate ACYW135(Men-C-ACYW135)** Date: \_\_\_\_\_  
**Possible side effects:** Soreness, swelling and/or redness at the injection site; fever (temperature of 38.5 C or higher); tiredness or drowsiness; irritability or crying; decreased appetite.
  
- Hepatitis B (HB)** Date: 1. \_\_\_\_\_ 2. \_\_\_\_\_ 3.\* \_\_\_\_\_  
**Possible side effects:** Soreness, swelling and/or redness at the injection site; fever (temperature of 38.5 C or higher); a small painless lump at the injection site that may take a few weeks to disappear but does not need any treatment; tiredness. \* A 2-dose schedule is used for individuals 11 – 15 years of age and a 3-dose schedule is used for all other age groups.
  
- Human Papillomavirus (HPV-9)** Date: 1. \_\_\_\_\_ 2. \_\_\_\_\_ 3.\* \_\_\_\_\_  
**Possible side effects:** Soreness, swelling and/or redness at the injection site; lump at injection site; fever (temperature of 38.5 C or higher); mild headache; nausea; diarrhea; tiredness; abdominal pain; dizziness; sore throat. \* A 2-dose schedule is used for individuals 11 – 15 years of age and a 3-dose schedule is used for all other age groups.
  
- Tetanus, diphtheria, acellular pertussis (Tdap)** Date: \_\_\_\_\_  
**Possible side effects:** Soreness, swelling and/or redness at the injection site; fever (temperature of 38.5 C or higher); mild headache.

**What to Do:**

- Give plenty of fluids to drink
- Dress your child in loose, light clothing
- If fever occurs, give fever reducing medication
- Apply a cool, damp cloth to the injection site

**More severe effects are extremely rare and if symptoms develop after you leave the clinic, call 9-1-1. After business hours, the Healthline can be reached at 811. It is important to always report serious or unexpected reactions to your health care provider.**

- Allergic reaction (hives; swelling of the face, lips or throat; wheezing; shock; difficulty breathing)
- Pain, or numbness and tingling in the hands and feet
- Convulsions or seizures
- Extreme drowsiness or unresponsiveness

Community/Public Health Nurse \_\_\_\_\_ Telephone \_\_\_\_\_

To keep a record of all your child’s immunizations please visit: <http://www.immunize.ca/en/app>